

Rec'd PGT/PTO 29 APR

10/533243

[Handwritten mark]

PA 1096598

REC'D 12 DEC 2003

WIPO

PC

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

November 21, 2003

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/422,861

FILING DATE: November 01, 2002

GB/03/4711

PRIORITY DOCUMENT

SUBMITTED OR TRANSMITTED IN COMPLIANCE WITH RULE 17.1(a) OR (b)

By Authority of the
COMMISSIONER OF PATENTS AND TRADEMARKS



[Handwritten signature of T. Lawrence]

T. LAWRENCE
Certifying Officer

BEST AVAILABLE COPY

PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION under 37 CFR 1.53 (c)

Docket Number 620-224

Type a plus sign (+) inside this box →

1/01/02



11059 S. PRO

INVENTOR(S)/APPLICANT(S)

LAST NAME
FREEMAN

FIRST NAME
Matthew

MIDDLE INITIAL

RESIDENCE (CITY AND EITHER STATE OR FOREIGN COUNTRY)
Cambridge, UNITED KINGDOM

TITLE OF THE INVENTION (280 characters)

ASSAYS, METHODS AND MEANS

CORRESPONDENCE ADDRESS

B. J. Sadoff
NIXON & VANDERHYE P C
1100 North Glebe Road
8th Floor
Arlington

STATE

Virginia

ZIP CODE

22201

COUNTRY

ENCLOSED APPLICATION PARTS (check all that apply)

☒ Specification

Number of Pages

38

☐ Applicant claims "small entity" status

☐ "Small entity" statement attached

☒ Drawing(s)

Number of Sheets

1

☐ Other (specify)

METHOD OF PAYMENT (check one)



A check or money order is enclosed to cover the Provisional filing fees (\$160.00)/(\$80.00)



The commissioner is hereby authorized to charge filing fees and credit

Deposit Account Number

14-1140

PROVISIONAL
FILING FEE
AMOUNT (\$)

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government



No.



Yes, the name of the U.S. Government agency and the Government contract number are:

Respectfully submitted,
SIGNATURE

DATE

November 1, 2002

TYPED or PRINTED NAME

B. J. Sadoff

REGISTRATION NO.
(if appropriate)

36



Additional inventors are being named on separately numbered sheets attached hereto

PROVISIONAL APPLICATION FILING ONLY

Burden Hour Statement This form is estimated to take 2 hours to complete. Time will vary depending upon the needs of the individual case. Any commitment of time you are required to complete this form should be sent to the Office of Assistance Quality and Enhancement Division, Patent and Trademark Office, Washington, DC 20231, and to the Office of Information and Regulatory Affairs, Office of Management and Budget (Project 0651-0037), Washington, DC 20231. NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Washington, DC 20231

Our Ref.: 620-224
NRS/CP6097463

U.S. PATENT APPLICATION

Inventor(s): Matthew FREEMAN

Invention: ASSAYS, METHODS AND MEANS

***NIXON & VANDERHYE P.C.
ATTORNEYS AT LAW
1100 NORTH GLEBE ROAD, 8TH FLOOR
ARLINGTON, VIRGINIA 22201-4714
(703) 816-4000
Facsimile (703) 816-4100***

SPECIFICATION



Assays, Methods and Means

The present invention relates to the invasive processes of protozoan pathogens and in particular to the provision of therapeutic compounds which block these processes and may thus be useful in treating protozoan pathogen infection.

Protozoan pathogens of the Apicomplexa family, which include the malaria parasite *P. falciparum*, express a number of membrane-tethered adhesion proteins that are essential for the recognition and binding of host (e.g. mammalian) cells. An essential step in the invasive processes by which these pathogens enter host cells is the release of these adhesion proteins from the surface of the pathogen by proteolytic cleavage. Some of these adhesion proteins are cleaved within the transmembrane domain (TMD) (Opitz et al (2002) EMBO J. 21 7 1577-1585).

The present inventors have discovered that the protease responsible for the cleavage of protozoan adhesion molecules is a member of the Rhomboid family. Rhomboid polypeptides are intra-membrane serine proteases, which are widely conserved throughout evolution and act on a range of physiological substrates. Inhibition of the protozoan Rhomboid activity reduces adhesion polypeptide cleavage and may thus reduce the invasiveness of the protozoan pathogen.

A first aspect of the invention provides a method for identifying and/or obtaining a compound which inhibits invasiveness of a protozoan pathogen, for example by inhibiting the activity of a protozoan Rhomboid polypeptide, which method comprises:

- (a) bringing into contact an isolated Rhomboid polypeptide and an isolated substrate polypeptide in the presence of a test compound; and
- (b) determining proteolytic cleavage of the substrate polypeptide.

The Rhomboid and substrate polypeptides may be contacted out under conditions in which the Rhomboid polypeptide normally catalyses proteolytic cleavage of the substrate polypeptide.

5

The polypeptides may be contacted in a reaction medium in an isolated form or may be comprised in a liposome or cell. The Rhomboid polypeptide may, for example, act on a membrane-bound substrate polypeptide to generate a soluble product, which is
10 detected.

Cleavage of the substrate polypeptide may be determined in the presence and absence of test compound. A reduction or decrease in cleavage in the presence of the test compound relative to the
15 absence of test compound may be indicative of the test compound being an inhibitor of protozoan Rhomboid protease activity. Such a compound may inhibit adhesive micronemal polypeptide cleavage and thus protozoan pathogen infectivity.

20 A Rhomboid polypeptide may be a protozoan Rhomboid protein, for example a Rhomboid polypeptide from an apicomplexan pathogen.

Amino acid residues of Rhomboid and substrate polypeptides are described in the present application with reference to their
25 position in the Drosophila Rhomboid-1 and Spitz sequences, respectively. It will be appreciated that the equivalent residues in other Rhomboid and substrate polypeptides may have a different position and number, because of differences in the amino acid sequence of each polypeptide. These differences may
30 occur, for example, through variations in the length of the N terminal domain. Equivalent residues in other Rhomboid and substrate polypeptides are easily recognisable by their overall sequence context and by their positions with respect to the TMDs.

35

A polypeptide which is a member of the Rhomboid family preferably comprises residues R152, G215, S217 and H281, more

preferably residues W151, R152, N169, G215, S217 and H281. The presence of these conserved residues may be used to identify Rhomboid polypeptides in other protozoan pathogens.

- 5 Preferably, a Rhomboid polypeptide comprises at least 4 TMDs, more preferably at least 5 TMDs, with residues N169, S217 and H281 each occurring in different TMD at about the same level in the lipid membrane bilayer.
- 10 Preferred Rhomboid polypeptides comprise a GxSx motif, for example GxSG.

A Rhomboid polypeptide may also comprise additional amino acid residues which are heterologous to the Rhomboid sequence. For
15 example, a Rhomboid polypeptide or fragment thereof may be included as part of a fusion protein, e.g. including a binding portion for a different ligand.

Whilst Rhomboid polypeptides may share relatively low overall
20 sequence identity, they may be reliably identified by bioinformatics techniques and manual inspection of key residues, as described herein.

A polypeptide which is a member of the Rhomboid family (i.e. a
25 rhomboid polypeptide) may be identified by the presence of a Rhomboid homology domain, as defined by the PFAM protein structure annotation project (Bateman A. et al (2000) The Pfam Protein Families Database Nucl. Acid. Res. 28 263-266). The Pfam rhomboid homology domain is built from a Hidden Markov Model
30 (HMM) using 83 rhomboid sequences as a seed. The Pfam 'rhomboid' domain has the pfam specific accession number PF01694.

Other methods suitable for use in identifying Rhomboid polypeptides are well known in the art.

35

Particularly valuable methods include the use of Hidden Markov Models built from groups of previously identified Rhomboid

proteins, including, but not limited to Drosophila Rhomboids 1-4. Such bio-informatics techniques are well known to those skilled in the art (Eddy S. R. Curr. Opin. Struct. Biol. 1996 6(3) 361-365).

5

Rhomboid polypeptides are preferably able to proteolytically cleave one or more of Drosophila proteins Spitz, Keren and Gurken but not similar type I transmembrane proteins like TGF α , Delta, EGFR, and TGN38.

10

Suitable substrate polypeptides are Type 1 membrane proteins with a single TMD orientated with an N terminal extracellular/luminal domain. A substrate polypeptide may comprise a transmembrane domain which has a lumen proximal region having one or more of the residues of residues 138-144 of the Drosophila Spitz sequence (ASIASGA), or a luminal region having an equivalent conformation, structure or three dimensional arrangement to that of residues 138-144 of the Drosophila Spitz sequence (ASIASGA).

20

In preferred embodiments, residues 1 to 7 of the TMD of a substrate polypeptide are not hydrophobic.

25

A substrate polypeptide may comprise a G residue within the portion of the TMD proximal to the luminal or extracellular domain of the polypeptide (i.e. between residues 1 and 8 of the TMD), preferably, the G residue being at a position equivalent to Spitz G143 (i.e the 6th residue of the TMD). Preferably, a substrate polypeptide comprises a GA or GG motif within the portion of the TMD proximal to the luminal or extracellular domain of the polypeptide (i.e. between residues 1 and 8 of the TMD), preferably at positions equivalent to G143 and A144 of the Spitz sequence (i.e the 6th and 7th residues of the TMD).

30

35

A substrate polypeptide may further comprise a triplet motif of small amino acid residues at positions equivalent to A138, S139 and I140 of Spitz (i.e. the 1st, 2nd and 3rd residues of the TMD

as numbered from the N terminal (extracellular) boundary of the TMD). Small residues include Gly, Ala, Ser, Ile, Leu, Val and Thr. Suitable motifs at this position include motifs which have Ala at a position equivalent to Spitz A138, such as AGG and ASI.

5 The presence of Phe at positions 1, 2 or 3 of the TMD of a polypeptide is a counter-indication for the polypeptide being a Rhomboid substrate (i.e. such a polypeptide is unlikely to be efficiently cleaved by a Rhomboid polypeptide).

10 Suitable substrate polypeptides may include protozoan adhesive micronemal polypeptides. An adhesive micronemal polypeptide may be from a protozoan pathogen, for example an Apicomplexan pathogen. Suitable adhesive micronemal polypeptides include transmembrane proteins that belong to the thrombospondin-
15 related-anonymous proteins (TRAPs) family.

A protozoan adhesive micronemal polypeptide which is a suitable substrate may include an AGG or AGL motif at residues 1, 2 and 3 of the TMD as described above and a GG motif at positions 6 and
20 7 of the TMD.

The protozoan pathogen may be a member of the Apicomplexa family, for example an apicomplexan pathogen selected from the group consisting of Plasmodium, Toxoplasma, Eimeria, Sarcocystis,
25 Babesia, Isospora, Cyclospora and Cryptosporidium, for example *P. falciparum*, *P. bergei*, *T. gondii*, *C. parvum*, *E. tenella*, *S. muris*, *Babesia bovis*, *Cyclospora belli*, *Theileria annulata* or *Theileria parva*.

30 Examples of suitable polypeptides include AMA1, MIC2, MIC6, MIC8, MIC12, (from *T. gondii*), AMA1, TRAP, CTRP (from *P. falciparum* and *P. berghei*), MIC1 and MIC4 (from *E. tenella*) or other polypeptide as shown in Table 2.

35 The substrate polypeptide and the Rhomboid polypeptide may each comprise an ER (endoplasmic reticulum) retention signal. For example, a rhomboid polypeptide may comprise a C terminal

(luminal) KDEL motif and a substrate may comprise a C terminal (cytoplasmic) KKXX motif (Jackson et al (1993) J. Cell Biol. 121(2) 317-333)

5 The substrate polypeptide may comprise a detectable label, such as green fluorescent protein (GFP), luciferase or alkaline phosphatase. This is preferably located in the soluble extracellular domain, allowing convenient detection of the soluble cleaved product and is particularly useful in automated
10 assays.

Methods for obtaining or identifying modulators, in particular inhibitors, of protozoan pathogen infectivity may be *in vivo* cell-based assays, or *in vitro* non-cell-based assays.

15 In *in vitro* assays, the rhomboid polypeptide and the substrate polypeptide may be isolated or contained in a liposome. Liposome based assays may be carried out using methods well-known in the art (Brenner C. et al (2000) Meths in Enzymol. 322 243-252,
20 Peters et al (2000) Biotechniques 28 1214-1219, Puglielli, H. and Hirschberg C. (1999) J. Biol. Chem. 274 35596-35600, Ramjeesingh, M. (1999) Meths in Enzymol. 294 227-246).

25 Preferably, assays according to the present invention take the form of *in vivo* assays. *In vivo* assays may be performed in a cell such as a protozoan cell (e.g. a Toxoplasma cell), a yeast strain, insect or mammalian cell line such as CHO, HeLa and COS cells, in which the relevant polypeptides or peptides are expressed from one or more vectors introduced into the cell.
30

In a preferred embodiment, the Rhomboid polypeptide and the substrate polypeptide may be expressed in a host cell from heterogeneous encoding nucleic acid.

35 Nucleic acid encoding the Rhomboid polypeptide and the substrate polypeptide may be contained on a single expression vector or on separate expression vectors.

Persons skilled in the art may vary the precise format of methods of the invention using routine skill and knowledge. It is not necessary to use the entire full-length proteins for

5 in vitro or in vivo assays of the invention. Polypeptide fragments as described herein which retain the activity of the full length protein may be generated and used in any suitable way known to those of skill in the art. Suitable ways of generating fragments include, but are not limited to,

10 recombinant expression of a fragment from encoding DNA. Such fragments may be generated by taking encoding DNA, identifying suitable restriction enzyme recognition sites either side of the portion to be expressed, and cutting out said portion from the DNA. The portion may then be operably linked to a suitable

15 promoter in a standard commercially available expression system. Another recombinant approach is to amplify the relevant portion of the DNA with suitable PCR primers. Small fragments (e.g. up to about 20 or 30 amino acids) may also be generated using peptide synthesis methods, which are well known in the art.

20 Another approach is to synthesise a nucleic acid comprising all or part of the coding sequence from a series of synthetic oligonucleotides. The coding sequence may be synthesised using a more optimal genetic code for the host cells (Kocken CH et al Infect Immun. (2002) 70(8):4471-6).

25 A Rhomboid polypeptide fragment may consist of fewer amino acid residues than said full-length polypeptide. Such a fragment may consist of 325 amino acids or less, 300 amino acids or less, or 275 amino acids or less and/or may consist of at least 100 amino acids, more preferably at least 150, 200, 250 or 300 amino

30 acids. A suitable fragment may comprise five TMDs.

Such a fragment preferably comprises residues equivalent to R152, G215, S217 and H281, more preferably residues W151, R152, N169, G215, S217 and H281 of the Drosophila Rhomboid-1 sequence (Acc No:P20350), which are important for the catalytic activity

35 of the protein and are highly conserved in the Rhomboid family.

A substrate polypeptide fragment comprises fewer residues than the full-length polypeptide and preferably comprises the transmembrane domain of the polypeptide. The TMD may be conveniently identified using commercially available software such as TMHMM (Krogh A. et al (2001) J. Mol. Biol. 305 567-580) and TmPred (Hofmann K & Stoffel W (1993) Biol Chem. Hoppe Seyler 374 166).

Combinatorial library technology (Schultz, JS (1996) Biotechnol. Prog. 12:729-743) provides an efficient way of testing a potentially vast number of different substances for ability to modulate activity of a polypeptide. Prior to or as well as being screened for modulation of activity, test substances may be screened for ability to interact with the Rhomboid polypeptide, e.g. in a yeast two-hybrid system (which requires that both the polypeptide and the test substance can be expressed in yeast from encoding nucleic acid). This may be used as a coarse screen prior to testing a substance for actual ability to modulate protease activity of the polypeptide.

The amount of test substance or compound, which may be added in a method of the invention, will normally be determined by trial and error depending upon the type of compound used. Typically, from about 0.01 to 100µM concentrations of putative inhibitor compound may be used, for example from 0.1 to 10µM.

Test compounds may be natural or synthetic chemical compounds used in drug screening programmes. Extracts of plants that contain several characterised or uncharacterised components may also be used.

Methods of the invention may comprise the step of identifying the test compound as an inhibitor of adhesive micronemal polypeptide cleavage.

One class of putative inhibitor compounds can be derived from the Rhomboid polypeptide and/or substrate polypeptide (e.g. adhesive micronemal polypeptide). Peptide fragments of from 5 to 40 amino acids, for example, from 6 to 10 amino acids may be tested for their ability to disrupt such interaction or activity.

The inhibitory properties of a peptide fragment as described above may be increased by the addition of one of the following groups to the C terminal: chloromethyl ketone, aldehyde and boronic acid. These groups are transition state analogues for serine, cysteine and threonine proteases. The N terminus of a peptide fragment may be blocked with carbobenzyl to inhibit aminopeptidases and improve stability (Proteolytic Enzymes 2nd Ed, Edited by R. Beynon and J. Bond Oxford University Press 2001).

Antibodies directed to the site of interaction in the protozoan Rhomboid polypeptide or adhesive micronemal protein form a further class of putative inhibitor compounds. Candidate inhibitor antibodies may be characterised and their binding regions determined to provide single chain antibodies and fragments thereof which are responsible for disrupting the interaction.

Antibodies may be obtained using techniques that are standard in the art. Methods of producing antibodies include immunising a mammal (e.g. mouse, rat, rabbit, horse, goat, sheep or monkey) with the protein or a fragment thereof. Antibodies may be obtained from immunised animals using any of a variety of techniques known in the art, and screened, preferably using binding of antibody to antigen of interest. For instance, Western blotting techniques or immunoprecipitation may be used (Armitage et al., 1992, Nature 357: 80-82). Isolation of antibodies and/or antibody-producing cells from an animal may be accompanied by a step of sacrificing the animal.

As an alternative or supplement to immunising a mammal with a peptide, an antibody specific for a protein may be obtained from a recombinantly produced library of expressed immunoglobulin variable domains, e.g. using lambda bacteriophage or filamentous bacteriophage which display functional immunoglobulin binding domains on their surfaces; for instance see WO92/01047. The library may be naive, that is constructed from sequences obtained from an organism which has not been immunised with any of the proteins (or fragments), or may be one constructed using sequences obtained from an organism which has been exposed to the antigen of interest.

Antibodies according to the present invention may be modified in a number of ways. Indeed, the term "antibody" should be construed as covering any binding substance having a binding domain with the required specificity. Thus the invention covers antibody fragments, derivatives, functional equivalents and homologues of antibodies, including synthetic molecules and molecules whose shape mimicks that of an antibody enabling it to bind an antigen or epitope.

Example antibody fragments, capable of binding an antigen or other binding partner are the Fab fragment consisting of the VL, VH, Cl and CH1 domains; the Fd fragment consisting of the VH and CH1 domains; the Fv fragment consisting of the VL and VH domains of a single arm of an antibody; the dAb fragment which consists of a VH domain; isolated CDR regions and F(ab')₂ fragments, a bivalent fragment including two Fab fragments linked by a disulphide bridge at the hinge region. Single chain Fv fragments are also included.

The reactivities of antibodies on a sample may be determined by any appropriate means. Tagging with individual reporter molecules is one possibility. The reporter molecules may directly or indirectly generate detectable, and preferably measurable, signals. The linkage of reporter molecules may be directly or indirectly, covalently, e.g. via a peptide bond or

non-covalently. Linkage via a peptide bond may be as a result of recombinant expression of a gene fusion that encodes antibody and reporter molecule. The mode of determining binding is not a feature of the present invention and those skilled in the art
 5 are able to choose a suitable mode according to their preference and general knowledge.

Antibodies may also be used in purifying and/or isolating Rhomboid and adhesive micronemal polypeptides for use in the
 10 present methods, for instance following production of the polypeptide or peptide by expression from encoding nucleic acid therefor.

Antibodies may be useful in a therapeutic context (which may include prophylaxis) to disrupt Rhomboid mediated cleavage of
 15 adhesive micronemal proteins and thus to reduce pathogen invasiveness in the treatment protozoan infections, including malaria.

20 Antibodies may be employed in accordance with the present invention for other therapeutic and non-therapeutic purposes which are discussed elsewhere herein.

Adhesive micronemal polypeptide cleavage is an essential step in
 25 the infection of host cells by protozoan pathogens. Compounds which inhibit adhesive micronemal polypeptide cleavage may thus inhibit protozoan pathogen invasiveness.

Methods of the invention may further comprise the step of
 30 determining the ability of said test compound to inhibit the invasiveness of a protozoan pathogen.

This may be achieved by contacting a host cell with a protozoan pathogen in the presence of the test compound under conditions
 35 in which the pathogen normally infects the host cell.

Invasiveness of a protozoan pathogen may be determined, for example, in the presence and absence of test compound, in tissue culture using hepatocytes or HepG2 cells. Alternatively, the ability of a sporozoite to glide on a glass slide may be
5 determined. This motility is tightly linked to invasiveness (Matuschewski K et al EMBO J (2002) 21 (7) 1597-1606).

In other assay systems, sporozoites may be injected into an animal model (e.g. rat) to which a test compound is administered
10 and the extent or amount of infection of hepatocytes within the animal determined, and compared to control animals (Matuschewski K et al supra).

A decrease or reduction in the rate of infection in the presence
15 relative to the absence of test compound is indicative that the test compound inhibits infectivity.

The test compound may further be isolated and/or manufactured/synthesised and subsequently formulated into a
20 pharmaceutical composition with a pharmaceutically acceptable excipient, vehicle or carrier.

It is desirable that compounds for use in therapeutic contexts preferentially or specifically inhibit protozoan Rhomboid
25 polypeptide relative to human Rhomboid polypeptide. Methods of the invention may include a further screen to identify such compounds.

A method may therefore comprise the further step of;
30 bringing into contact an isolated human Rhomboid polypeptide and a polypeptide substrate in the presence of the test compound; and, determining proteolytic cleavage of the substrate by the human Rhomboid polypeptide.

35 A human Rhomboid polypeptide may be selected from the group consisting of Human RHBDL-1 (Human Rhomboid-1: Pascall and Brown

(1998) FEBS Lett. 429, 337-340), Human RHBDL-2 (NM_017821) and Human RHBDL-3.

5 Suitable polypeptide substrates for human Rhomboids are described above and are cleaved by the Rhomboid polypeptide within the transmembrane domain.

10 A suitable substrate polypeptide may comprise a transmembrane motif which has one or more residues of the Drosophila Spitz ASIASGA motif within the region proximal to the luminal or extracellular domain of the polypeptide (i.e. residues 1 to 8 of the TMD starting at the luminal boundary) or may comprise a transmembrane motif which none of the residues of the Drosophila Spitz ASIASGA motif, but which instead possess a motif having an
15 equivalent structure which is cleaved by Rhomboid polypeptide (e.g. Gurken or other sequence shown in Figure 1). Preferably a substrate polypeptide comprises one or both of motifs described above in the region proximal to the luminal or extracellular domain of the polypeptide.

20 Following identification of a compound using a method of the invention described herein, a method may further comprise modifying the compound to optimise the pharmaceutical properties thereof.

25 The modification of a 'lead' compound identified as biologically active is a known approach to the development of pharmaceuticals and may be desirable where the active compound is difficult or expensive to synthesise or where it is unsuitable for a
30 particular method of administration, e.g. peptides are not well suited as active agents for oral compositions as they tend to be quickly degraded by proteases in the alimentary canal. Modification of a known active compound (for example, to produce a mimetic) may be used to avoid randomly screening large number
35 of molecules for a target property.

Modification of a 'lead' compound to optimise its pharmaceutical properties commonly comprises several steps. Firstly, the particular parts of the compound that are critical and/or important in determining the target property are determined. In the case of a peptide, this can be done by systematically varying the amino acid residues in the peptide, e.g. by substituting each residue in turn. These parts or residues constituting the active region of the compound are known as its "pharmacophore".

10

Once the pharmacophore has been found, its structure is modelled according to its physical properties, e.g. stereochemistry, bonding, size and/or charge, using data from a range of sources, e.g. spectroscopic techniques, X-ray diffraction data and NMR.

15

Computational analysis, similarity mapping (which models the charge and/or volume of a pharmacophore, rather than the bonding between atoms) and other techniques can be used in this modelling process.

20

In a variant of this approach, the three-dimensional structure of the ligand and its binding partner are modelled. This can be especially useful where the ligand and/or binding partner change conformation on binding, allowing the model to take account of this in the optimisation of the lead compound.

25

A template molecule is then selected onto which chemical groups that mimic the pharmacophore can be grafted. The template molecule and the chemical groups grafted on to it can conveniently be selected so that the modified compound is easy to synthesise, is likely to be pharmacologically acceptable, and does not degrade *in vivo*, while retaining the biological activity of the lead compound. The modified compounds found by this approach can then be screened to see whether they have the target property, or to what extent they exhibit it. Modified compounds include mimetics of the lead compound.

30

35

Further optimisation or modification can then be carried out to arrive at one or more final compounds for *in vivo* or clinical testing.

- 5 Rhomboid and adhesive miconemal polypeptides may also be used in methods of designing mimetics which are suitable for inhibiting protozoan pathogen infectivity.

10 The present invention provides a method of designing mimetics having the biological activity of inhibiting the Rhomboid mediated cleavage of adhesive miconemal polypeptides and thus inhibiting protozoan pathogen invasiveness, said method comprising:

- 15 (i) analysing a compound having the biological activity to determine the amino acid residues essential and important for the activity to define a pharmacophore; and,
(ii) modelling the pharmacophore to design and/or screen candidate mimetics having the biological activity.

- 20 A suitable compound may be, for example, a protozoan Rhomboid polypeptide or fragment as described herein.

25 Suitable modelling techniques are known in the art. This includes the design of so-called "mimetics" which involves the study of the functional interactions of the molecules and the design of compounds which contain functional groups arranged in such a manner that they could reproduced those interactions.

30 The modelling and modification of a 'lead' compound to optimise its properties, including the production of mimetics, is described above.

35 The activity or function of a protozoan Rhomboid polypeptide may be inhibited, as noted, by means of a compound that interferes in some way with the interaction of protozoan Rhomboid with adhesive micronemal polypeptides or other suitable substrate polypeptides. An alternative approach to inhibition employs

regulation at the nucleic acid level to inhibit activity or function by down-regulating production of protozoan Rhomboid.

For instance, expression of a gene may be inhibited using anti-sense or RNAi technology. The use of these approaches to
5 down-regulate gene expression is now well-established in the art.

Anti-sense oligonucleotides may be designed to hybridise to the complementary sequence of nucleic acid, pre-mRNA or mature mRNA, interfering with the production of Rhomboid polypeptide so that
10 its expression is reduced or completely or substantially completely prevented. In addition to targeting coding sequence, antisense techniques may be used to target control sequences of a gene, e.g. in the 5' flanking sequence, whereby the antisense oligonucleotides can interfere with expression control
15 sequences. The construction of antisense sequences and their use is described for example in Peyman and Ulman, Chemical Reviews, 90:543-584, (1990) and Crooke, Ann. Rev. Pharmacol. Toxicol., 32:329-376, (1992).

Oligonucleotides may be generated *in vitro* or *ex vivo* for
20 administration or anti-sense RNA may be generated *in vivo* within cells in which down-regulation is desired. Thus, double-stranded DNA may be placed under the control of a promoter in a "reverse orientation" such that transcription of the anti-sense strand of the DNA yields RNA which is complementary to normal mRNA
25 transcribed from the sense strand of the target gene. The complementary anti-sense RNA sequence is thought then to bind with mRNA to form a duplex, inhibiting translation of the endogenous mRNA from the target gene into protein. Whether or not this is the actual mode of action is still uncertain.
30 However, it is established fact that the technique works.

The complete sequence corresponding to the coding sequence in reverse orientation need not be used. For example fragments of sufficient length may be used. It is a routine matter for the person skilled in the art to screen fragments of various sizes

and from various parts of the coding or flanking sequences of a gene to optimise the level of anti-sense inhibition. It may be advantageous to include the initiating methionine ATG codon, and perhaps one or more nucleotides upstream of the initiating
5 codon. A suitable fragment may have about 14-23 nucleotides, e.g. about 15, 16 or 17.

An alternative to anti-sense is to use a copy of all or part of the target gene inserted in sense, that is the same, orientation as the target gene, to achieve reduction in expression of the
10 target gene by co-suppression; Angell & Baulcombe (1997) The EMBO Journal 16,12:3675-3684; and Voinnet & Baulcombe (1997) Nature 389: pg 553). Double stranded RNA (dsRNA) has been found to be even more effective in gene silencing than both sense or antisense strands alone (Fire A. et al Nature, Vol 391, (1998))
15 and has been used effectively in Plasmodium (Malhotra P et al Mol. Microb. (2002) 45(5) 1245-1254; McRobert L et al Mol. Biochem. Parasitol. (2002) 119(2) 273-278). dsRNA mediated silencing is gene specific and is often termed RNA interference (RNAi).

20 RNA interference is a two step process. First, dsRNA is cleaved within the cell to yield short interfering RNAs (siRNAs) of about 21-23nt length with 5' terminal phosphate and 3' short overhangs (~2nt). The siRNAs target the corresponding mRNA sequence specifically for destruction (Zamore P.D. Nature
25 Structural Biology, 8, 9, 746-750; (2001)

RNAi may be also be efficiently induced using chemically synthesized siRNA duplexes of the same structure with 3'-overhang ends (Zamore PD et al Cell, 101, 25-33, (2000)). Synthetic siRNA duplexes have been shown to specifically
30 suppress expression of endogenous and heterologous genes in a wide range of mammalian cell lines (Elbashir SM. et al. Nature, 411, 494-498, (2001)).

Another possibility is that nucleic acid is used which on transcription produces a ribozyme, able to cut nucleic acid at a

specific site - thus also useful in influencing gene expression. Background references for ribozymes include Kashani-Sabet and Scanlon, 1995, *Cancer Gene Therapy*, 2(3): 213-223, and Mercola and Cohen, 1995, *Cancer Gene Therapy*, 2(1), 47-59.

5

Thus, a modulator of protozoan Rhomboid activity and thus a modulator of protozoan pathogen infectivity may comprise a nucleic acid molecule comprising all or part of a Rhomboid coding sequence shown in Table 1 or the complement thereof

10

Such a molecule may suppress the expression of protozoan Rhomboid polypeptide and may comprise a sense or anti-sense Rhomboid coding sequence or may be an Rhomboid specific ribozyme, according to the type of suppression to be employed.

15

The type of suppression will also determine whether the molecule is double or single stranded and whether it is RNA or DNA.

20

Another aspect of the present invention provides a method of producing a pharmaceutical composition comprising; identifying a compound which inhibits the invasiveness of a protozoan pathogen using a method described herein; and, admixing the compound identified thereby with a pharmaceutically acceptable carrier.

25

As described above, a method of the invention may comprise the step of modifying the compound to optimise the pharmaceutical properties thereof.

30

Another aspect of the invention provides a method for preparing a pharmaceutical composition for treating a protozoan pathogen infection comprising;

35

i) identifying a compound which modulates the proteolytic activity of a Rhomboid polypeptide,
ii) synthesising the identified compound, and;
iii) incorporating the compound into a pharmaceutical composition.

The identified compound may be synthesised using conventional chemical synthesis methodologies. Methods for the development and optimisation of synthetic routes are well known to a skilled person.

The compound may be modified and/or optimised as described above.

Incorporating the compound into a pharmaceutical composition may include admixing the synthesised compound with a pharmaceutically acceptable carrier or excipient.

Another aspect of the invention provides a compound which modulates protozoan pathogen infectivity obtained by a method as described herein. Such a compound may comprise or consist of a peptide fragment of a protozoan Rhomboid polypeptide.

Another aspect of the invention provides a pharmaceutical composition comprising a compound which modulates the proteolytic activity of a Rhomboid polypeptide obtained by a method described herein.

In other aspects the invention provides the use of a compound obtained by a method described herein in the manufacture of a composition for treatment of a protozoan pathogen infection and a method comprising administration of a composition obtained by a method described herein to a patient for treatment of a protozoan pathogen infection.

Protozoan pathogens are described above. Disorders associated with infection with a protozoan pathogen include malaria, toxoplasmosis, cryptosporidiosis, diarrhoea associated with *Isospora belli* or *Cyclospora cayetanensis* infection, and various livestock disorders associated with *Eimeria* infection.

Whether it is a polypeptide, antibody, peptide, anti-sense, sense or siRNA nucleic acid molecule, small molecule, mimetic or other pharmaceutically useful compound according to the present invention that is to be given to an individual, administration is preferably in a "prophylactically effective amount" or a "therapeutically effective amount" (as the case may be, although prophylaxis may be considered therapy), this being sufficient to show benefit to the individual. The actual amount administered, and rate and time-course of administration, will depend on the nature and severity of what is being treated. Prescription of treatment, e.g. decisions on dosage etc, is within the responsibility of general practitioners and other medical doctors.

A composition may be administered alone or in combination with other treatments, either simultaneously or sequentially dependent upon the condition to be treated.

Pharmaceutical compositions according to the present invention, and for use in accordance with the present invention, may include, in addition to active ingredient, a pharmaceutically acceptable excipient, carrier, buffer, stabiliser or other materials well known to those skilled in the art. Such materials should be non-toxic and should not interfere with the efficacy of the active ingredient. The precise nature of the carrier or other material will depend on the route of administration, which may be oral, or by injection, e.g. cutaneous, subcutaneous or intravenous.

Pharmaceutical compositions for oral administration may be in tablet, capsule, powder or liquid form. A tablet may include a solid carrier such as gelatin or an adjuvant.¹ Liquid pharmaceutical compositions generally include a liquid carrier such as water, petroleum, animal or vegetable oils, mineral oil or synthetic oil. Physiological saline solution, dextrose or other saccharide solution or glycols such as ethylene glycol, propylene glycol or polyethylene glycol may be included.

For intravenous, cutaneous or subcutaneous injection, or injection at the site of affliction, the active ingredient will be in the form of a parenterally acceptable aqueous solution which is pyrogen-free and has suitable pH, isotonicity and stability. Those of relevant skill in the art are well able to prepare suitable solutions using, for example, isotonic vehicles such as Sodium Chloride Injection, Ringer's Injection, Lactated Ringer's Injection. Preservatives, stabilisers, buffers, antioxidants and/or other additives may be included, as required.

Liposomes, particularly cationic liposomes, may be used in carrier formulations.

Examples of techniques and protocols mentioned above can be found in Remington's Pharmaceutical Sciences, 16th edition, Osol, A. (ed), 1980.

Another aspect of the invention provides a method of identifying a protozoan Rhomboid polypeptide comprising,
(a) providing a test Rhomboid polypeptide,
(b) bringing into contact a substrate polypeptide and the test Rhomboid polypeptide under conditions in which the substrate polypeptide is normally proteolytically cleaved; and,
(c) determining cleavage of the substrate polypeptide.

A suitable test Rhomboid polypeptide may comprise an amino acid sequence encoded by a nucleic acid sequence shown in Table 1.

Screening databases may identify other suitable test Rhomboid polypeptides, in particular, databases of protozoan nucleic acid sequence, using the bioinformatics techniques discussed above.

Suitable substrate polypeptides include protozoan adhesive micronemal polypeptides. A suitable adhesive micronemal polypeptide may belong to the thrombospondin-related-anonymous proteins (TRAPs) family. Examples of suitable polypeptides

include AMA1, MIC2, MIC6, MIC8, MIC12, (from *T. gondii*), TRAP, CTRP (from *P. falciparum* and *P. berghei*), MIC1 and MIC4 (from *E. tenella*) or fragments thereof which comprise the transmembrane domain.

5

Other aspects of the invention relate to protozoan Rhomboid polypeptides and encoding nucleic acids.

An aspect of the invention provides a protozoan Rhomboid polypeptide which proteolytically cleaves the transmembrane domain of an adhesive micronemal polypeptide.

10

Such a polypeptide may have a sequence encoded by a nucleic acid sequence shown in Table 1 or may be a fragment of such a sequence which consists of fewer residues than the full-length Rhomboid polypeptide.

15

The KDEL ER retention sequence is not found in natural Rhomboid polypeptides and directs the expressed Rhomboid polypeptide to be retained the ER (endoplasmic reticulum) rather than the miconemes of the protozoan cell. As described below, Rhomboid polypeptides labelled with an ER retention sequence such as KDEL may be particularly useful in certain embodiments of the invention, as proteolytic cleavage by such polypeptides avoids potential problems with variations in secretion efficiency.

20

An aspect of the invention thus provides an isolated Rhomboid polypeptide as described above comprising an C terminal ER retention sequence. A suitable retention sequence consists of the amino acid sequence KDEL.

25

Another aspect of the invention provides an isolated nucleic acid encoding a Rhomboid polypeptide as described above.

30

The coding sequence may be a nucleic acid sequence listed in Table 1 or it may be a mutant, variant, derivative or allele of a sequence listed. The sequence may differ from a sequence of

35

Table 1 by a change, which is one or more of addition,
insertion, deletion and substitution of one or more nucleotides
of the sequence shown. Changes to a nucleotide sequence may
result in an amino acid change at the protein level, or not, as
5 determined by the genetic code.

Thus, nucleic acid according to the present invention may
include a sequence different from a sequence listed in Table 1
yet encode a polypeptide with the same amino acid sequence. As
10 described above, codon usage may be adjusted in order to express
an amino acid sequence in a particular host system, such as a
mammalian cell.

An isolated nucleic acid may share greater than about 10%
15 sequence identity with a nucleic acid sequence of Table 1
greater than 20%, greater than 30%, greater than 40%, greater
than 50%, greater than 60%, greater than about 70%, greater than
about 80%, greater than about 90%, or greater than about 95%.

20 The present invention also extends to nucleic acid that
hybridizes with a sequence listed in Table 1 under stringent
conditions. Suitable conditions include, e.g. for sequences
that are about 80-90% identical; hybridisation overnight at 42°C
in 0.25M Na₂HPO₄, pH 7.2, 6.5% SDS, 10% dextran sulfate and a
25 final wash at 55°C in 0.1 X SSC, 0.1% SDS. For sequences that
are greater than about 90% identical, suitable conditions
include hybridization overnight at 65°C in 0.25M Na₂HPO₄, pH 7.2,
6.5% SDS, 10% dextran sulfate and a final wash at 60°C in 0.1X
SSC, 0.1% SDS.

30 A convenient way of producing a polypeptide for use in assays
and methods according to the present invention is to express
nucleic acid encoding it, by use of the nucleic acid in an
expression system. Accordingly, the present invention also
35 encompasses a method of making a polypeptide (as disclosed), the
method including expression from nucleic acid encoding the
polypeptide and testing for Rhomboid protease activity. This

may conveniently be achieved by growing a host cell in culture, containing such a vector, under appropriate conditions which cause or allow expression of the polypeptide. Polypeptides may also be expressed in *in vitro* systems, such as reticulocyte
5 lysate.

Another aspect of the present invention therefore provides a method of producing a Rhomboid polypeptide comprising:

- 10 (a) causing expression from nucleic acid which encodes a protozoan Rhomboid polypeptide in a suitable expression system to produce the polypeptide recombinantly;
- (b) testing the recombinantly produced polypeptide for Rhomboid protease activity.

15 Suitable nucleic acid sequences include a nucleic acid sequence encoding a protozoan rhomboid polypeptide a mutant, variant or allele thereof as described herein.

A polypeptide may be isolated and/or purified (e.g. using an
20 antibody) for instance after production by expression from encoding nucleic acid (for which see below). Thus, a polypeptide may be provided free or substantially free from contaminants with which it is naturally associated (if it is a naturally-occurring polypeptide). A polypeptide may be provided
25 free or substantially free of other polypeptides.

The recombinantly produced polypeptide may be isolated and/or tested for Rhomboid protease activity by determination of the cleavage of a substrate polypeptide such as an adhesive
30 micronemal polypeptide upon incubation of the polypeptide with the substrate polypeptide.

An isolated nucleic acid as described herein, for example a nucleic acid encoding a protozoan Rhomboid polypeptide, may be
35 comprised in a vector. Such a vector may further include a nucleic acid sequence encoding a substrate polypeptide such as a protozoan adhesive micronemal protein. Suitable vectors can be

chosen or constructed, containing appropriate regulatory sequences, including promoter sequences, terminator fragments, polyadenylation sequences, enhancer sequences, marker genes and other sequences as appropriate. Vectors may be plasmids, viral
5 e.g. 'phage, or phagemid, as appropriate. For further details see, for example, Molecular Cloning: a Laboratory Manual: 2nd edition, Sambrook et al., 1989, Cold Spring Harbor Laboratory Press. Many known techniques and protocols for manipulation of nucleic acid, for example in preparation of nucleic acid
10 constructs, mutagenesis, sequencing, introduction of DNA into cells and gene expression, and analysis of proteins, are described in detail in Current Protocols in Molecular Biology, Ausubel et al. eds., John Wiley & Sons, 1992.

15 Systems for cloning and expression of a polypeptide in a variety of different host cells are well known. Suitable host cells include bacteria, eukaryotic cells such as mammalian and yeast, baculovirus and protozoan systems. Mammalian cell lines available in the art for expression of a heterologous
20 polypeptide include Chinese hamster ovary cells, HeLa cells, baby hamster kidney cells, COS cells and many others. A common, preferred bacterial host is *E. coli*.

Further aspects of the invention provide a host cell containing
25 heterologous nucleic acid encoding a protozoan Rhomboid polypeptide, including a Rhomboid polypeptide which has a KDEL tag or which is a fragment of a full length Rhomboid sequence, and a host cell containing heterologous nucleic acid encoding a protozoan Rhomboid polypeptide and a substrate polypeptide, for
30 example a protozoan adhesive micronemal polypeptide. Nucleic acid encoding the protozoan Rhomboid polypeptide and substrate polypeptide may be present on a single nucleic acid construct or vector within the host cell or nucleic acid encoding the two polypeptides may be present on separate constructs or vectors.

35

The nucleic acid may be integrated into the genome (e.g. chromosome) of the host cell. Integration may be promoted by

inclusion of sequences that promote recombination with the genome, in accordance with standard techniques. The nucleic acid may be on an extra-chromosomal vector within the cell.

5 The introduction of nucleic acid into a host cell, which may (particularly for *in vitro* introduction) be generally referred to without limitation as "transformation", may employ any available technique. For eukaryotic cells, suitable techniques may include calcium phosphate transfection, DEAE-Dextran,
10 electroporation, liposome-mediated transfection and transduction using retrovirus or other virus, e.g. vaccinia or, for insect cells, baculovirus. For bacterial cells, suitable techniques may include calcium chloride transformation, electroporation and transfection using bacteriophage.

15 Marker genes such as antibiotic resistance or sensitivity genes may be used in identifying clones containing nucleic acid of interest, as is well known in the art.

20 The introduction may be followed by causing or allowing expression from the nucleic acid, e.g. by culturing host cells (which may include cells actually transformed although more likely the cells will be descendants of the transformed cells) under conditions for expression of the gene, so that the encoded
25 polypeptide is produced.

A Rhomboid polypeptide may be co-expressed in a host cell with a substrate polypeptide, such as a protozoan adhesive micronemal polypeptide, and the Rhomboid serine protease activity
30 determined by determining cleavage of the substrate polypeptide. Cleavage may be determined by determining the presence or absence of soluble cleavage products which may be secreted into the culture medium.

35 Aspects of the present invention will now be illustrated with reference to the following experimental exemplification, by way of example and not limitation. Further aspects and embodiments

will be apparent to those of ordinary skill in the art. All documents mentioned in this specification are hereby incorporated herein by reference.

5 Figure 1 shows an alignment of transmembrane domains of micronemal proteins from apicomplexan species with *Drosophila* Spitz and Keren. All are single-pass type 1 transmembrane proteins. Black lines indicate the predicted TMD regions: the line above the sequences predicts the micronemal protein TMDs; 10 the line below predicts the TMDs of Spitz and Keren. The GA or GG motif approximately seven residues into the TMD (double underline) and the conserved small residues near the luminal/extracellular face of the TMD (single underline) are essential for rhomboid cleavage of Spitz and are underscored. 15 Note also the conserved tyrosine (Y) at the cytoplasmic face of the TMDs. Accession numbers of the sequences are shown in Table 2.

Table 1a shows examples of *Plasmodium falciparum* rhomboids as 20 predicted by the Pfam motif-finding algorithm, with a score of greater than 10. Four annotations are listed: sequencing consortium annotation (Sanger), plus 3 automatic gene prediction algorithms, FullPhat, GlimmerM and Genefinder. The approximate position on the chromosome is also provided along with the 25 predicted sequence around the rhomboid active site. Rhomboids 2 and 5 are predicted to be mitochondrial, based on the existence of a mitochondrial targeting sequence predicted by MitoProt or Predotar. These genes represent the most reliable prediction of true rhomboids in the published *P. falciparum* 30 genome sequence as predicted by the Pfam motif-finding algorithm, with a score of greater than 10.

Table 1b shows an example of a candidate *Toxoplasma gondii* rhomboid (ref: <http://ParaDB.cis.upenn.edu/toxo/index.html>)

35 Table 2 shows examples of substrates for rhomboid proteases from apicomplexan species.

Experimental

Identification of Plasmodium Rhomboid polypeptides.

Rhomboid polypeptides were identified in the published
 5 Plasmodium genome sequence by using PFAM as described above,
 selecting scores higher than 10, followed by visual inspection
 to identify key Rhomboid residues.

A number of putative Rhomboids were identified using PFAM. These
 10 were further analysed for the presence of residues equivalent to
 residues N169, G215, S217 and H281 of Drosophila Rhomboid-1,
 which are required for catalytic activity. Candidates identified
 as being within the subclass of mitochondrial rhomboids were
 also excluded by prediction of the existence of a mitochondrial
 15 targeting sequence with PREDOTAR algorithm
 (www.inra.fr/predotar) or MitoProt (Claros M., et al Eur. J.
 Biochem. (1996) 241 779-786)

Rhomboid polypeptides identified by this approach are listed in
 20 Table 1.

Cloning of Plasmodium Rhomboid

Rhomboid coding sequences are amplified by PCR from Plasmodium
 genomic DNA (where introns are absent) or from cDNA (where
 25 introns are present) using the protocols set out in Sambrook &
 Russell Molecular Cloning, A Laboratory Manual, 3rd Edition, Cold
 Spring Harbor Laboratory Press, 2001, and Ausubel et al, Short
 Protocols in Molecular Biology, John Wiley and Sons, 1992.

Codon usage may be adjusted as described in Kocken et al (2002)
 30 Infect. Immun. 70(8) 4471-4476. Standard PCR mutagenesis
 techniques may be used to replace particular AT rich regions of
 the coding sequence or the entire coding sequence may be
 replaced using synthetic techniques. Cloned sequences may be
 35 epitope tagged to facilitate immunological detection of the
 expressed polypeptide.

The cloned sequences are then inserted into a pCDNA3.1 (Invitrogen) to express them in cell culture.

Cloning of adhesive micronemal proteins of Plasmodium

5 Oligonucleotide primers for amplification of the adhesive micronemal proteins of Plasmodium, as listed in Table 2 are designed based on the published sequences using conventional primer design software.

10 The coding sequences are amplified using PCR, and cloned into a standard mammalian expression vector for expression in cell culture. Expression is detected using available anti-TRAP antibodies.

15 Determining Rhomboid Activity

Rhomboid polypeptides are co-expressed with each of the adhesive micronemal proteins as follows;

COS cells are grown in DMEM medium (supplemented with 10% foetal calf serum), and transfected with FUGENE 6 Transfection Reagent (Roche).

20 Cells are co-transfected in 35mm culture wells with 25-250ng of;
(a) Rhomboid construct comprising a protozoan Rhomboid coding sequence inserted into the vector pCDNA 3.1+ (Invitrogen), and;
25 (b) Substrate construct containing the adhesive micronemal coding sequence inserted into pCDNA 3.1+(Invitrogen).

Empty plasmid is used to bring the total DNA to 1µg per well.

30 Transfection of construct (b) into COS cells in the absence of construct(a) acts as a control for endogenous cleavage of the substrate. Optionally, a metalloprotease inhibitor such as batimastat may be used to minimize endogenous substrate cleavage in the assay.

24-30 hours post-transfection the medium is replaced with serum-free medium; this is harvested 24 hours later and cells are lysed in SDS-sample buffer.

5 For some experiments the serum-free medium may be supplemented with the metalloprotease inhibitor batimastat (British Biotech) or ilomostat (Calbiochem). For inhibitor assays, a test compound may be included in the serum-free medium.

10 Supernatants are collected from the wells and assayed for the presence of GFP in the medium using conventional Western Blotting techniques.

For example, GFP may be captured with a polyclonal or monoclonal
15 antibody (Santa Cruz Biologicals), washed and then the captured GFP detected with a polyclonal or monoclonal antibody conjugated to an enzyme (capture ELISA) or with a fluorescent label.

For ELISA, a suitable polyclonal anti-GFP conjugated to horse-
20 radish peroxidase or to alkaline phosphatase may be used. Such a conjugate is preferred since the number of incubations required is reduced. Alternatively a biotinylated anti-GFP antibody in combination with an avidin or streptavidin enzyme conjugate could be used.

25 For a fluorescence assay, Europium- or Terbium-labelled antibody or streptavidin are suitable (e.g. Delphia or Lance reagents, Perkin Elmer). These are labels with a long fluorescence lifetime and can improve the signal:noise ratio.

30 A variation of the above is to replace GFP in the GFP-adhesive protein construct (or to add to the construct) with an enzyme label at the N-terminus to give a direct assay for the cleaved substrate in the medium. Suitable enzymes include Renilla
35 luciferase (Lui, J., and Escher, A. (1999) Gene 237, 153-159) and secreted alkaline phosphatase sequence (SEAP) (Clontech).

Demonstrate in vivo role of rhomboid in infectivity of parasites

Mutant strains of *P. falciparum* are generated in which the candidate rhomboid protein is knocked out. The infectiousness of the mutant parasite is then determined.

5

The knockout is done either by standard targeted gene disruption (Sultan et al Cell 90 511) or, more conveniently, by RNA interference (RNAi) as described below.

- 10 100 µg of RNA corresponding to each candidate Rhomboid gene is synthesized by *in vitro* transcription from 5 µg linearized plasmid templates according to manufacturer's instructions (Promega Ribomax system). The resulting RNA is purified using the RNeasy protocol (Qiagen), denatured by boiling, and annealed
- 15 in 1mM Tris-HCl pH7.4, 1mM EDTA overnight. The resulting dsRNA is then added to a culture of *P. falciparum* sporozoites (Malhotra et al Mol. Microb. 45 1245-1254).

- The reduced infectivity or gliding behaviour of the knockout parasites relative to the wild-type controls confirms that
- 20 rhomboid regulates the infectious process.

- Impairment of the proteolysis of the adhesive micronemal proteins may also be established in the rhomboid mutant (or
- 25 RNAi) cells, by determining the production of cleaved product by Western blotting using anti-TRAP antibodies (Sharma et al Infection & Immunity 64 2172-2179, Gantt et al Infection & Immunity 68 3667-3673).

30

Sanger	FullPhat	GlimmerM	Genefinder	Chromosome	Approx location	Motif	Rhomboid
PFE03040c	chr5.phat_92	chr5.glm_74	chr5.gen_274	5	289kb	GSSG	Rho1
PFE0755c	chr5.phat_172	chr5.glm_161	chr5.gen_236	5	624kb	GASG	Rho2 (mito)
PF11_0150	chr11.phat_164	chr11.glm_163	Not predicted	11	538kb	GAST	Rho3
PF13_0312	chr13.phat_576	chr13.glm_633	chr13.gen_395	13	2.22Mb	GSSS	Rho5 (mito)
PF13_0241	chr13.phat_452	chr13.glm_494	chr13.gen_456	13	1.73Mb	GASG	Rho4
PF14_0110	chr14.phat_108	Not predicted	chr14.gen_705	14	453kb	SSSS	Rho6 (?)
MAL8P1.16	chr8.phat_47	chr8.glm_34	chr8.gen_290	8	135kb	GAST	Rho7

Table 1a

Toxoplasma gondii consensus ID	Genbank accession	Motif	Rhomboid
Ctoxogual 3360	AA531635	GAST	TgRho1

Table 1b

Protein	Species	Accession no.
PfTRAP	Plasmodium falciparum	U67764
PfCTRP	Plasmodium falciparum	U34363
PfAMA1	Plasmodium falciparum	AF277003
EBL-175	Plasmodium falciparum	AAM33518
MAEBL	Plasmodium falciparum	AY042084
TgMIC2	Toxoplasma gondii	TGU62660
TgMIC6	Toxoplasma gondii	AF110270
TgMIC12	Toxoplasma gondii	Opitz et al.
TgAMA1	Toxoplasma gondii	AF010264
PbTRAP	Plasmodium berghei	U67763
EtMIC1	Eimeria tenella	AF032905
EtMIC4	Eimeria tenella	CAC34726
Sm70	Sarcocystis muris	AAK35069

Table 2

Claims

1. An method for identifying and/or obtaining a compound which inhibits infectivity of a protozoan pathogen, which method
5 comprises:
 - (a) bringing into contact an isolated Rhomboid polypeptide and an isolated substrate polypeptide in the presence of a test compound; and
 - (b) determining proteolytic cleavage of the substrate protein.
- 10 2. A method according to claim 1 wherein the protozoan pathogen is an apicomplexan pathogen.
3. A method according to claim 2 wherein the apicomplexan
15 pathogen is selected from the group consisting of Plasmodium, Toxoplasma, Eimeria, Sarcocystis, Cyclospora, Isospora, Cryptosporidium, Babesia and Theileria.
4. A method according to any one of the preceding claims
20 wherein the Rhomboid polypeptide is a protozoan Rhomboid protein
5. A method according to claim 4 wherein the Rhomboid polypeptide is encoded by a nucleic acid sequence shown in Table
1.
- 25 6. A method according to any one of the preceding claims wherein the substrate polypeptide comprises a luminal domain and a TMD, the TMD having a region proximal to the luminal domain which comprises one or more of residues 138-144 of the
30 Drosophila Spitz sequence (ASIASGA).
7. A method according to claim 6 wherein the substrate polypeptide comprises a TMD and a luminal domain, the TMD having a region proximal to a luminal domain which has the sequence of
35 residues 138-144 of Drosophila Spitz.

8. A method according to claim 6 wherein the substrate polypeptide is an adhesive micronemal polypeptide.

5 9. An assay method according to claim 8 wherein the substrate polypeptide is encoded by a nucleic acid sequence shown in Table 2.

10 10. A method according to any one of the preceding claims wherein the substrate polypeptide and the Rhomboid polypeptide comprise ER (endoplasmic reticulum) retention signals.

11. A method according to claim 10 wherein the endoplasmic reticulum retention signals are KDEL or KKXX.

15 12. A method according to any one of the preceding claims wherein the substrate polypeptide comprises an extracellular domain having a detectable label.

20 13. A method according to any one of the preceding claims wherein said Rhomboid polypeptide and said substrate polypeptide are expressed in a host cell from heterogeneous nucleic acid.

14. A method according to any one of the preceding claims comprising the further step of;

25 (a) bringing into contact an isolated human Rhomboid polypeptide and a polypeptide substrate in the presence of a test compound; and,

(b) determining proteolytic cleavage of the substrate by the human Rhomboid polypeptide.

30

15. A method according to any one of the preceding claims comprising identifying said test compound as a modulator of adhesive micronemal polypeptide cleavage.

35 16. A method according to claim 15 further comprising determining the ability of said test compound to inhibit the invasiveness of a protozoan pathogen.

17. A method according to claim 15 or claim 16 comprising isolating said test compound.
- 5 18. A method according to claim 17 comprising formulating said test compound in a pharmaceutical composition with a pharmaceutically acceptable excipient, vehicle or carrier.
- 10 19. A compound which modulates protozoan pathogen infectivity obtained by a method of any one of claims 1 to 16.
20. A compound according to claim 19 comprising a peptide fragment of a protozoan Rhomboid polypeptide.
- 15 21. A method of producing a pharmaceutical composition comprising;
identifying a compound which inhibits the infectivity of a protozoan pathogen using a method according to any one of claims 1 to 16; and,
20 admixing the compound identified thereby with a pharmaceutically acceptable carrier.
- 25 22. A method according to claim 21 comprising the step of modifying the compound to optimise the pharmaceutical properties thereof.
- 30 23. A method for preparing a pharmaceutical composition for treating a protozoan pathogen infection comprising;
i) identifying a compound which modulates the proteolytic activity of a Rhomboid polypeptide,
ii) synthesising the identified compound, and,
iii) incorporating the compound into a pharmaceutical composition.
- 35 24. A pharmaceutical composition comprising a compound according to claim 19 or claim 20.

25. Use of a compound according to claim 19 or claim 20 in the manufacture of a composition for treatment of a protozoan pathogen infection.

5 26. A method comprising administration of a composition according to claim 19 or claim 20 to a patient for treatment of a protozoan pathogen infection.

10 27. A method according to claim 26 wherein the protozoan pathogen is an apicomplexan pathogen selected from the group consisting of Plasmodium, Babesia, Theileria, Toxoplasma, Eimeria, Sarcocystis, Cyclospora, Isospora and Cryptosporidium.

15 28. A method of identifying a protozoan Rhomboid polypeptide comprising;
 (a) providing a test protozoan Rhomboid polypeptide,
 (b) bringing into contact a substrate polypeptide and the test Rhomboid polypeptide under conditions in which the substrate polypeptide is normally proteolytically cleaved; and,
 20 (c) determining cleavage of the substrate polypeptide.

25 29. A method according to claim 28 wherein the test Rhomboid polypeptide comprises an amino acid sequence encoded by a nucleic acid sequence shown in Table 1.

30. A method according to claim 28 or claim 29 wherein the substrate polypeptide comprises the luminal region of the TMD of Spitz, Gurken or Keren.

30 31. A method according to claim 28 or claim 29 wherein the substrate polypeptide comprises an amino acid sequence encoded by a nucleic acid sequence shown in Table 2.

ABSTRACTAssays, Methods and Means

5 The present invention relates to the recognition that intra-
membrane proteases of the Rhomboid family play an important role
in the invasive processes of protozoan pathogens. Inhibition of
these Rhomboid proteases may thus be useful in treating
protozoan pathogen infection and methods and means relating to
10 this inhibition are provided.

5	EtMIC4	...GFPTAAV <u>AGGV</u> <u>GGV</u> LLLLAAVGGGV--AAAY
	TgMIC2	...SGIAGAI <u>AGGV</u> <u>IGGL</u> ILLGAAG---GASY
	TgMIC6	...SGHAGAI <u>AGGV</u> <u>IGGL</u> LLLLSAAGAG--VAY
	TgMIC12	...GVPVAAI <u>AGGV</u> <u>VGGV</u> LLLIAGGAGA--AVY
	PbTRAP	...SNNGIKI <u>AGGI</u> <u>IGGL</u> AIIGCIG---VGY
10	PfTRAP	...SDNKYKI <u>AGGI</u> <u>AGGL</u> ALLACAG---LAY
	EtMIC1	...GFPTAAV <u>AGGV</u> <u>AGGV</u> LAIAAGAG---AFY
	Sm70	...GMPTAAI <u>AGGI</u> <u>VGGV</u> LLLLGAAGGG--AAAY
	TgAMA1	...GSNTALI <u>AGLA</u> <u>VGGV</u> LLLLALLGG---GCY
	DmSPITZ	...PRPMLEK <u>ASIAS</u> <u>GAM</u> CALVFMLFVCLAFYL
15	DmKEREN	...NRVMLEK <u>ASIV</u> <u>SGAT</u> LALLFMAMCCVVL

lumenal/extracellular

cytoplasmic

Figure 1

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.